Artificial intelligence is used in a smart city to understand the emotional pulse of its residents

M. Praveena¹, K. Sirisha², G. Siva Prasad³, Dr. D Jebakumar immanuel⁴, U. Sai Bhavya⁵

^{1, 2, 3, 4, 5} Department of Computer Science and Engineering,

^{1, 2, 3, 5}QIS College of Engineering and Technology, Ongole, Andhra Pradesh, India

⁴SNS College of Engineering, Coimbatore 641107, Tamilnadu

praveena mulle@qiscet.edu.in¹, sirisha .k@qiscet.edu.in², g.sivaprasad@qiscet.edu.in³, jebakumarimmanuel@snsce.ac.in⁴, saibhavya.u@qiscet.edu.in⁵

Corresponding Author Mail: qispublications@qiscet.edu.in

Article Info	Abstract: India, which is in the tropical wet and dry area, receives a huge								
Page Number: 50-60	amount of precipitation each year, with the monsoon season being the								
Publication Issue:	main contributor. The environment resulting challenges also has led to								
Vol 69 No. 1 (2020)	significant improvements in the average rain distribution and its volatility,								
	as well as the severity and frequency of severe rainfalls. Rain also exhibits								
	great temporal and geographic volatility. Different Time-series								
	prognostication models for predicting rain, including Holt's Linear Trend								
	methodology (HLTM), Generalized Auto-Regressive Conditional								
	Heteroskedasticity (GARCH), Holt's Winter Seasonal Methodology								
	(HWSM), and seasonal Autoregressive Integrated Moving Average, were								
	used in this study (SARIMA). In order to analyse the most basic time-								
	series prognostication model, a comparison analysis was built. The								
	HWSM model was determined to have the lowest error rate when								
	compared to the other models. Mean square error (MSE), Root Mean								
	square error (RMSE), and Mean Absolute Error are the analysis criteria								
Article Received: 15 September	used to compare different time-series rain forecast methods (MAE). The								
2020	HWSM model exhibited, by far, the lowest error rate of the other models,								
Revised: 24 October 2020	with error rates of 4.767 and 4.343 RMSE, 1.51 and 1.432 MAE, and								
Accepted: 26 November Publication: 30 December 2020	25.65 and 24.75 MSE for datasets 1 and 2, respectively.								

Introduction

Since rainfall data should be recorded over a period of time, it is considered a statistical analysis. In areas including business, ecological prognostication, and technological prognostication, statistical prognostication is employed as a decision-support tool. Due to connectivity, a variety of models and methodologies for statistical forecasting have been developed that enable various types of inputs, anticipated outcomes, and straightforward implementation. Our country's economy is based primarily on agriculture. The value of the harvest has changed more recently due to uncertain environmental condition patterns and other market changes. Despite this, farmers continue to ignore these hazards, which results in destroyed crops and significant losses. Because of this, they are unable to determine which harvest would result in higher revenues.

Lack of understanding about various agricultural pests and resulting yearly precipitation amounts causes farms to be destroyed. The economy of Asia still consists mostly of agriculture, which would be strongly correlated with the amount of precipitation received annually [1]. Geophysical and atmospheric science both depend on accurate precipitation forecasts. On the other side, current systems frequently provide low error rates and perform well in terms of prediction. Numerous situations call for poor performance from mathematical prediction models. The whole prediction performance and error rate for precipitation prediction were investigated throughout this examination, which examined and assessed a number of time-series foretelling models. A variety of techniques are used to estimate the annual precipitation, including Multi-Layer Perceptron (MLP), Deep Neural Networks (DNN), Random Call Forest (RDF), K-Nearest Neighbors (KNN), K-Suggests that clump (KMC), Support Vector Machine (SVM), Simulated Neural Network (SNN), Transfer learning (TL), automobile Encoder Neural Network (AENN), and Convolution Neural Network (CNN) [Such methods also incorporate the techniques from the metric capacity unit [3] and Deep Learning (DL) [4], both of which have application potential. The following might serve as a summary of the rest of this article: In Section Two, pertinent analysis is presented, in Section Three, the study's region is covered, and in Section Four, several approaches for the prediction model for precipitation estimation are presented, Section five contains the observations and outcomes, and Section vi concludes the analysis.

1. Related Works

Researchers, academicians, and specialists used a variety of techniques to predict the frequency of precipitation by taking into consideration monumental efficiency assessment metrics, specifically RMSE, variance (SD), Learning rate (LR), MAE, Accuracy, MSE, Mean Absolute proportion Error (MAPE), F1-Measure, Sensitivity, and Nash-Sutcliffe efficiency (NSE) (PET). An overview of precipitation prediction models is provided in Table 1.

Resear ch	The appr oach empl oyed to predi ct rainf all	Meteoro logical characte ristics used	Rainf all predi ction befor e based on Locat ion and no. of days	Eval uatio n desi gns for meas urin g perf orma nce	Dataset	Experiment al Survey	Limitations
--------------	--	---	--	--	---------	-------------------------	-------------

 Table 1. Rainfall Prediction Models – A Survey.

A.B. Wicaks ono Putra et al., (2020). [5]	CNN , SNN	Sunlight radiation , Pressure on sea level	Indon esia &Thi rty days	MSE , MA PE	Samarinda meteorolog ical centers from 2006 to 2019	MSE=0.00 098 MAPE=0.0 005	Improve accuracy by using AENN & DNN models
A.R. Naik et al., (2020). [6]	SVM , decis ion tree, RDF	moisture , warmth, and Baromet ric Pressure	Globa l surve y	Ada m opti mize r	It is a survey DNN predi accurately	paper. They co ct rainfall	ncluded that predictions
M.M.R . Khan et al., (2020). [7]	Long Short Term Mem ory	Tempera ture and geopote ntial altitude	Bangl adesh 7 days	Stan dard devi ation and Mea n error	Kaggle website [13] for 1901-2015	Mean = 0.38 Std Devi- 0.63	By acceleratin g processing, failure rate can be reduced
E.Huss ein et al (2020). [8]	SVM	Sunlight radiation , pressure in sea, Breeze speed	USA & thirty days	MA E and RM SE	Kaggle website - American dataset	Mapping rainfall, forecasting weather with minimum errors	Globally extended
A Kumar et al., (2021). [9]	ML desig ns	Pressure in sea, cloud point	India & 7 days	Acc urac y	Rainfall in summer from 1991- 2019	Accuracy= 89.98%	More parameters included for research
Y.R.Sa ri et al., (2020). [10]	CNN & 1 day	Tempera ture, breeze speed	Indon esia & 2 week	Acc urac y, miss	Dataset from reference [14]	Accuracy= 81.46% & miss = 0.0018.	Suggestion s for improving CNN

							2320-980.
		and	S	ing			
		cloud					
S. Hariba bu et al., (2021). [11]	SVM , MLP , KM C	wetness, warm	Not specif ied	Acc urac y	Data from reference [12]	High precision levels generated	Efficiency can be improved for other factors.

2. Case Study Location

Telangana, India's 29th state, was established on June 2, 2014, following the detachment from Andhra Pradesam (AP), situated in south India. AP surmised geographic directions of 15°55' N and 19°55' N scope and 77°10' E to 81°50' E longitude, and has limits with AP. The long stretch of Telangana state's South West Rainstorm (SWM) precipitation in August (31.35%), sought after by 19.6%, 30.6%, 18.4 percent for June, July, and September months. The state encounters annual rainfall of 78.65% only from SWM season.

Fig.1 portrays the regions of Telangana in India. Fig.2 portrays state's divisions. Territories in Telangana's north-east regions considered the most water compared with the territory's of south during June to September in SWM season.

Table2 displays the greatest storm over the most recent thirty years (1989-2018) for each SWM seasons in Telangana state.



Fig. 1: Regions of Telangana Location



Fig. 2: state divisions

Table 2.	The ma	aximum	rainfall	in	each	SW	/M	season

Month	Maximum	downpour	experienced	in	Year
	111111				
June	234.8				2000
July	492.6				2013
August	426.5				1990
September	248.7				2007
June- September	1133.1				2013
Annual	1384.7				2013

3. Methodology

Figure 3 illustrates the evaluation and comparison of SARIMA, GARCH, HWSM, and HLTM as part of the methodology used to forecast the amount of precipitation.



Fig. 3: Working Principle

3.1. Data Selection and Preprocessing

The two datasets utilized are DataSet-1 (DS-1), which comprises normalized data on atmospheric phenomena for each district and months from 1951 to 2000, and DataSet-2 (DS-2), which contains mean atmospheric phenomena data for each district from 1901 to 2015. These datasets have been compiled by IMD [12]. MAE, RMSE and MAE were utilized to evaluate the classification model's dependability.

3.2. SARIMA

The philosophy used to anticipate the precipitation include the Informational collection Determination, Information Preprocessing, and assessing and contrasting GARCH, SARIMA, HLTM and HWSM which are displayed in Fig.3.

3.3. GARCH

GARCH models are applied when the measure of variability of the error term is either constant or uniform. So, in a mathematical model based on statistical assumptions, heteroskedasticity refers to the uneven pattern of parameter variation. Data from time series are subjected to these analysis, which examines the contingent measure of variability. When applied to datasets having a positive excess kurtosis error and a large SD, it quantifies all multi-dimensional relationship. The model is a modified version of ARCH that additionally comprises of MA.

3.4. HWSM

Holt Winter's Exponential Smoothing and triple exponential smoothing are some names for it. This time-series forecasting methodology projects parameter estimates by accounting for both trend and seasonality.

In order to apply HWSM, periodic variables must first be exponentially smoothed. Level and trend are also included. Over a time overarching pattern of values is used frequently. In

the sequence, the levels are the average values. Seasonality, commonly referred to as a repetition of a particular set of recordings taken with specific breaks. It consists of a evaluation formula and 3 smoothing formulae: Le_i, Tr_i, and Pe_i, which are expressed in eqn. (1), eqn. (2), eqn. (3) & eqn. (4) with 3 smoothing parameters α , β , and γ .

This formula shows the adjusted mean value of non-seasonal prediction and the season calibrated observation for period i. The trend formula is equivalent to HLTM. An identified mean value of the current time interval indicator and the previous season is shown by the below periodic formula.

Level Formula	$Le_i = \alpha(a_i - Pe_{i-l}) + (1 - \alpha)(Le_{i-1} + Tr_{i-1})$	eqn. (1)
Trend Formula (2)	$Tr_i = \beta(Le_i - Le_{i-1}) + (1 - \beta)Tr_{i-1}$	eqn.
Seasonality/Periodicity Forr	nula $Pe_i = \gamma(a_i - Le_i) + (1 - \gamma)Pe_{i-1}$	eqn. (3)

Prediction formula	$F_{i+n} = Le_i + n Tr_i + Pe_{i+n-1}$	eqn. (4)

Were Le represents Level, F indicates prediction at n duration, a represents observation, Tr is trend parameter, Pe specifies time interval indicator, α , β , and γ are constants that must be evaluated similar to MSE model as small as possible, 1 gives the duration of time interval patterns for α , β and γ , where α , β and γ resides between 0 and 1.

3.5. HLTM

Enhanced SES enables forecasting of data with a trend. The SES method known as HLTM is combined with the level and trend. Now this requires 3 formulas to be justified algebraically. The HLTM is composed of two smoothing equations, Le_i and Tr_i, which are represented by Equations (5), (6), and (7), respectively, and have smoothing parameters of and. α and β .

Level Formula	$Le_i = \alpha(a_i - Pe_{i-1}) + (1 - \alpha)(Le_{i-1} + Tr_{i-1})$	(5)
Trend Formula	$Tr_i = \beta(Le_i - Le_{i-1}) + (1 - \beta)Tr_{i-1}$	(6)
Prediction formula	$F_{i+n} = Le_i + n Tr_i$	(7)

4. Results and Discussion

We evaluated these approaches using their RMSE, MAE, and MSE values. Table 3 presents a comparison of the various Time-Series forecasting methods. Table 3 shows that HWSM had the lowest error rates for DS-1 and DS-2, respectively, 4.767 and 4.343 RMSE, 1.51 and 1.432 MAE, and 27.65 and 24.75 MSE. Fig. 4 and Fig. 5 shows the representation of MAE, MSE and RMSE values for the DS-1, DS-2 for each of the models of time series forecasting that were taken into consideration.



Fig. 4: Forecasting models of MAE, MSE and RMSE values of Dataset - 1



Fig. 5: Forecasting models of MAE, MSE and RMSE values of Dataset - 2

Model	RMSE		MAE		MSE	
	DS-1	DS-2	DS-1	DS-2	DS-1	DS-2
SARIMA	6.856	6.706	3.1	2.541	30.260	28.542
GARCH	6.682	5.280	3.678	3.59	49.021	39.443
HWSM	4.767	4.343	1.51	1.432	27.65	24.75
HLTM	7.912	8.241	3.462	3.394	55.723	52.390

Table 3. Comparison Forecasting Models in time intervals.

5. Conclusion

This research offered a Comparative Forecasting Models in time intervals for evaluating rainfall in Telangana state. The results show that HWSM outperforms the alternative strategies in terms of RMSE, MSE and MAE. The HLTM, HWSM, SARIMA, and GARCH

time-series forecasting models were evaluated. It is clear that HWSM has the lowest error rates for datasets 1 and 2, with 4.767 and 4.343 RMSE, 1.51 and 4.432 MAE, and 27.65 and 24.75 MSE, respectively.

References

- Sunori S K, Mittal A, Juneja P, Prakash P O, Alagh R and Maurya S 2021 Prediction of rainfall using GRNN and neurofuzzy techniques Proc. Int. Conf. on Advancement in Technology 2021 1-7.
- Hong K and Kang T 2021 A study on rainfall prediction based on meteorological time series Proc. Int. Conf. on Ubiquitous and Future Networks 2021 302-304.
- 3. Ponnamperuma N and Rajapakse L 2021 Comparison of time series forecast models for rainfall and drought prediction Proc. Int. Conf. on Moratuwa Engineering Research **2021** 626-631.
- Kaliyaperumal K, Rahim A, Sharma D K, Regin R, Vashisht S and Phasinam K 2021 Rainfall prediction using deep mining strategy for detection Proc. Int. Conf. on Smart Electronics and Communication 2021 1623-28.
- Putra A B, Malani R, Suprapty B and Gaffar A F 2020 A deep auto encoder semi convolution neural network for yearly rainfall prediction Proc. Int. Conf. on Intelligent Technology and Its Applications 2020 205-210.
- Naik A R, Deorankar A V and Ambhore P B 2020 Rainfall prediction based on deep neural network: a review Proc. Int. Conf. on Innovative Mechanisms for Industry Applications 2020 98-101.
- Khan M M, Siddique M A, Sakib S, Aziz A, Tasawar I K and Hossain Z 2020 Prediction of temperature and rainfall in bangladesh using long short term memory recurrent neural networks Proc. Int. Conf. on Multidisciplinary Studies and Innovative Technologies 2020 1- 6.
- Hussein E, Ghaziasgar M and Thron C 2020 Regional rainfall prediction using support vector machine classification of large-scale precipitation maps Proc. Int. Conf. on Information Fusion 2020 1-8.
- Sari Y R, Djamal E C and Nugraha F 2020 Daily rainfall prediction using one dimensional convolutional neural networks Proc. Int. Conf. on Computer and Informatics Engineering 2020 90-95.
- Haribabu S, Gupta G S, Kumar P N and Rajendran P S 2021 Prediction of flood by rainfall using MLP classifier of neural network model Proc. Int. Conf. on Communication and Electronics Systems 2021 1360-65.
- 11. Pal AB, Mishra PK. Trend analysis of rainfall, temperature and runoff data: a case study of Rangoon watershed in Nepal. International Journal of Student's Research in Technology & Management. 2017;2321–2543,5(3);21–38.
- Naveena K, Elzopy AK, Chaturvedi KA, Chandran MK, Gopinath G, Surendran U. Trend analysis of long-term rainfall temperature data for Ethopia. South African Geographical Journal; 2020. Available:https://doi.org/10.1080/03736245 .2020.1835699
- 13. Singh O, Arya P, Chaudhary BS. On rising temperature trends at Dehradun in Doon

valley of Uttarakhand, India. Journal of Earth System Science. 2013;122:613–622.

- 14. Zhang X, Vincent LA, Wd H, Niitsoo A. Temperature and rainfall trends in Canada during the 20th Century. AtmosphereOcean. 2000;38(3):395-429.
- 15. Fulekar M, Kale. Impact of climate change: Indian scenario. University News. 2010; 48(24):14-20.
- Mohammad P, Goswami A. Temperature and precipitation trend over 139 major Indian cities: an assessment over a century. Modeling Earth Systems and Environment. 2019;5:1481–1493 Available:https://doi.org/10.1007/s40808-019-00642-7
- 17. Gajbhiye S, Meshram C, Singh SK, Srivastava PK, Islam T. Precipitation trend analysis of Sindh River basin, India, from 102-year record (1901-2002). Atmospheric Science Letters. 2016;17:71–77.
- Kumar R, Gautam HR. Climate change and its impact on agricultural productivity in India. Journal of Climatology and Weather Forecasting. 2014;2:109. Available:https://doi.org/10.4172/2332- 2594. 1000109
- 19. Goswami BN. Increasing trend of extreme rain events over India in a warming
environment.Science.2016;314,1442-1445.Available:https://doi.org/10.1126/science.1132027
- 20. Dash SK, Jenamani RK, Kalsi SR, Panda SK. Some evidence of climate change in twentieth-century India. Climate Change. 2007;85:299–321.
- 21. Vittal B, Reddy KM. Statistical Analysis of rainfall moving trend in Telangana State. International Journal of Scientific Research in Mathematical and Statistical Sciences. 2019;6(4):08-15.
- 22. Navatha N, Sreenivas G, Umareddy R. Rainfall and temperature trends in Jagtial district of Telangana state. International Journal of Environment and Climate Change. 2021;11(11):47-59.
- 23. Jyothy AS, Murthy SD, Mallikarjuna P. Climate change impacts on seasonal rainfall trends in the regions of Andhra Pradesh and Telangana States, India. Journal of The Institution of Engineers (India). 2021;102(3):673-685.
- 24. Bhavana CP, Munirajappa R, Surendra HS, Rathod S. Modeling of daily rainfall using Gamma probability distribution. Jyostna et al.; IJECC, 12(11): 3405-3413, 2022; Article no.IJECC.93271 3413 Environment and Ecology. 2012;30(3B): 884-887.
- 25. Praveen B, Talukdar S, Shahfahad, Mahato S, Mondal J, Pritee S, Islam TMdR, Rahman A. Analyzing trend and forecasting of rainfall changes in India using non-parametrical and machine learning approaches. Nature Journal. 2020;10:10342
- 26. P Ramprakash, M Sakthivadivel, N Krishnaraj, J Ramprasath. "Host-based Intrusion Detection System using Sequence of System Calls" International Journal of Engineering and Management Research, Vandana Publications, Volume 4, Issue 2, 241-247, 2014
- 27. N Krishnaraj, S Smys."A multihoming ACO-MDV routing for maximum power efficiency in an IoT environment" Wireless Personal Communications 109 (1), 243-256, 2019.
- N Krishnaraj, R Bhuvanesh Kumar, D Rajeshwar, T Sanjay Kumar, Implementation of energy aware modified distance vector routing protocol for energy efficiency in wireless sensor networks, 2020 International Conference on Inventive Computation Technologies (ICICT),201-204

- Ibrahim, S. Jafar Ali, and M. Thangamani. "Enhanced singular value decomposition for prediction of drugs and diseases with hepatocellular carcinoma based on multi-source bat algorithm based random walk." Measurement 141 (2019): 176-183. https://doi.org/10.1016/j.measurement.2019.02.056
- 30. Ibrahim, Jafar Ali S., S. Rajasekar, Varsha, M. Karunakaran, K. Kasirajan, Kalyan NS Chakravarthy, V. Kumar, and K. J. Kaur. "Recent advances in performance and effect of Zr doping with ZnO thin film sensor in ammonia vapour sensing." GLOBAL NEST JOURNAL 23, no. 4 (2021): 526-531. https://doi.org/10.30955/gnj.004020 , https://journal.gnest.org/publication/gnest_04020
- N.S. Kalyan Chakravarthy, B. Karthikeyan, K. Alhaf Malik, D.Bujji Babbu, K. Nithya S.Jafar Ali Ibrahim, Survey of Cooperative Routing Algorithms in Wireless Sensor Networks, Journal of Annals of the Romanian Society for Cell Biology ,5316-5320, 2021
- 32. Rajmohan, G, Chinnappan, CV, John William, AD, Chandrakrishan Balakrishnan, S, Anand Muthu, B, Manogaran, G. Revamping land coverage analysis using aerial satellite image mapping. Trans Emerging Tel Tech. 2021; 32:e3927. https://doi.org/10.1002/ett.3927
- 33. Vignesh, C.C., Sivaparthipan, C.B., Daniel, J.A. et al. Adjacent Node based Energetic Association Factor Routing Protocol in Wireless Sensor Networks. Wireless Pers Commun 119, 3255–3270 (2021). https://doi.org/10.1007/s11277-021-08397-0.
- C Chandru Vignesh, S Karthik, Predicting the position of adjacent nodes with QoS in mobile ad hoc networks, Journal of Multimedia Tools and Applications, Springer US,Vol 79, 8445-8457,2020